



# Effect of Crude Extracts of Six Plants on Vegetative Growth of Soybean (*Glycine max*Merr.)

Zulfa Zakiah<sup>1</sup>, Irfan Suliansyah<sup>2</sup>, Amri Bakhtiar<sup>3</sup>, Mansyurdin<sup>4\*</sup>

<sup>1</sup>Department of Biology, Faculty of Mathematics and Natural Sciences, Tanjungpura University, Pontianak, Indonesia, [zulfazakiah@gmail.com](mailto:zulfazakiah@gmail.com)

<sup>2</sup>Department of Agronomy, Faculty of Agriculture, Andalas University, Padang, Indonesia, [Irfan.suliansyah@yahoo.com](mailto:Irfan.suliansyah@yahoo.com)

<sup>3</sup>Department of Biological Pharmacy, Faculty of Pharmacy, Andalas University, Padang, Indonesia, [amribakhtiar@gmail.com](mailto:amribakhtiar@gmail.com)

<sup>4</sup>Department of Biology, Faculty of Mathematics and Natural Sciences, Andalas University, Padang, Indonesia, [mansyurdin@gmail.com](mailto:mansyurdin@gmail.com)

\*Corresponding Author

---

**Abstract:** Boosting production by applying natural products derived from readily available plant sources to increase vegetative growth of soybean could help ensure food sovereignty, security and affordability of this important source of dietary protein. A preliminary trial using crude extracts from six potential growth stimulating sources suggested while treatment with *C. asiatica*, *M. esculenta* (cassava) and *G. linearis* leaf extract increased soybean plant height and leaf area, *A. scholaris* stem bark, *G. mangostana* pericarp and *Zea mays* silk reduced growth. Promising crude extracts from the first trial were retested at five dilute concentrations on 14-day old soybean plants using nested design with four replications. Analysis of variance and Duncan's Multiple Range Test ( $p \leq 0.05$ ) showed that crude extracts of cassava leaves, *C. asiatica* and *G. linearis* increased plant height, while only *C. asiatica* increased leaf area. The highest plant height and widest leaf area were obtained by applying 25mg/l crude extract of *C. asiatica*.

**Keywords:** soybean, vegetative growth, crude extract, growth stimulating.

---

## Introduction

A green revolution program using chemical fertilizers does not contribute to productivity enhancements after a certain saturation point. An alternative solution for increasing the growth of soybeans can be done using growth promoting substances from natural products. Up until now, the plant growth promoting properties of secondary metabolites of higher plants have been studied and used on a limited number of crops. Some plant crude extracts have, in fact, been reported to increase the plant growth. An extract of garlic, aloe and henna increased the fresh weight of *Schefflera arboricola* (Hanafy *et al.*,



**2012).** The application of red grape skin extract and alfalfa hydrolysate to pepper plants resulted in increased leaf biomass and weight of green peppers while increasing the growth and number of red peppers (**Ertani *et al.*, 2015**). Treating maize (*Zea mays* L.) with an extract derived from red grape skin (*Vitis vinifera* L.), blueberry fruit (*Vaccinium vitis-idaea* L.) and hawthorn leaves (*Crataegus monogina* Jacq.) resulted in increased root and leaf biomass, chlorophyll and sugar content (**Ertani *et al.*, 2016**). The application of *Moringa oleifera* leaf extract increased growth and yield of tomato (**Culver *et al.*, 2012**), maize and common beans (**Mvumi *et al.*, 2012**), some growth parameters and photosynthesis rate of rocket (*Eruca vesicaria* subsp. *sativa*) (**Abdalla, 2014**), germination and growth of wheat (Sarmin, 2014) and yield of sesame (*Sesamum indicum*) (**Muhamman and Mohammed, 2014**).

Aqueous extract of arjun tree leaves (*Terminalia arjuna*) increased germination and growth of turnip and okra. Treatment of aqueous extract of *T. belerica* increased germination and growth of spinach, while aqueous extract of neem (*Azadiracta indica*) inhibited a percentage of germination (**Talukder *et al.*, 2015**).

Secondary metabolite compounds including terpenoids, flavonoids, alkaloids and phenolic compounds have a potential as stimulants or inhibitors of plant growth. Secondary metabolites of some plants have been reported to have a wide range of bioactivity. *Centella asiatica* contains triterpenoids (terpenoids), flavonoids, and steroids (**Biradar and Rachetti, 2013; James and Dubery, 2011; Singh *et al.*, 2012**). Corn (*Zea mays*) silk contains steroids, flavonoids, and phenolic compounds (**Hasanudin *et al.*, 2012; Bhaigyabati *et al.*, 2011**). Cassava leaves contain flavonoids, terpenoids, alkaloids, steroids and saponins and phenolic compounds (**Siyumbawe *et al.*, 2014; Anbuselvi and Balamurugan, 2014; Nuwamanya *et al.*, 2014**). *Gleichenia linearis* fern leaves contain flavonoids and kaempferol (**Yusuf *et al.*, 2003; Jubahar, *et al.*, 2006; Dey, 2011**). The pericarp of mangosteen (*Garcinia mangostana*) fruit contains flavonoids, phenolic compounds and xanthones (**Zarena and Sakar, 2012**). Stem bark of *Alstonia scholaris* contains alkaloids, flavonoids, steroids, saponins and phenolic compounds (**Pankti *et al.*, 2012; Pratap *et al.*, 2013**).



Many of these plants that have been discovered to contain secondary metabolites are readily available in a tropical context. However, as far as the author is aware, none have been tested on the growth of soybeans (*Glycine max* Merr.). This study investigates the effects of secondary metabolites of some commonly available plants on the growth of this strategic crop.

## MATERIALS AND METHODS

### Preparation of crude extracts

Extracts from six plants were used in this research: cassava leaves (*M. esculenta*), *C. asiatica* stems and leaves, *G. linearis* leaves, corn silk, *A. scholaris* stem bark, and the pericarp of mangosteen fruit. Crude extracts of plants were prepared by maceration and boiling. Dry powder of *C. asiatica*, *A. scholaris* stem bark and pericarp of mangosteen fruits were macerated in methanol 70% (2 days), the extracts were filtered and concentrated in a rotary evaporator (45°C) and the final result was a solid crude extract (**Demissie and Lele, 2013**). Fresh cassava leaves and *G. linearis* were boiled in water (2 hours) then pressed and the extracts were filtered. After 3 days the crude extract, a yellow sediment was collected (**Jubahar, et al., 2006**).

### Phytochemical Analysis

The secondary metabolite contents in the crude extracts were analyzed using phytochemical analysis. For analyzing flavonoids, crude extract was diluted in 5 ml methanol and some drops of water, 10 mg of Mg powder, and two drops of HCl 5N added. A red color indicates the presence of flavonoids (**Shahid-Ud-Duaula and Anwarul, 2009**).

For analyzing steroids/terpenoids the Liebermann–Burchard reaction was used, crude extract was diluted in 5 ml chloroform and 5 ml water. The chloroform mixture/solution was passed through a small column containing active charcoal and then collected on a plate. After drying, 2 drops of anhydride acetate and 2 drops H<sub>2</sub>SO<sub>4</sub> were added, formation of a reddish brown ring confirms the presence of terpenes (**Obianime and Uche, 2008**). A Change of colour from violet to blue confirms the presence of steroids (**Boxi et al., 2010**).



For analyzing polar/semipolar phenolic compounds and saponin, crude extract was diluted in 5ml chloroform and 5ml water. Then one drop of the aqueous phase was placed on a plate and mixed with 2 drops of 1% FeCl<sub>3</sub> for polar phenolic compounds and mixed with some drops of methanol and 2 drops of 1% FeCl<sub>3</sub> for semipolar phenolic compounds. A red coloured solution with acetic acid indicates the presence of tannins and phenolic compounds (**Boxi *et al.*, 2010**).

For analyzing alkaloid, crude extract was diluted in 5ml chloroform ammonia 0.05N and 5ml water. 5-10 drops of H<sub>2</sub>SO<sub>4</sub> 2N was added to the chloroform phase and shaken slowly, then the acid phase was transferred into a reaction tube and 2 drops of Meyer's reagent were added (**Adegoke *et al.*, 2010**).

For saponin, some drops of the aqueous phase were transferred into a reaction tube and shaken quickly (1 minute). If the foam was stable up to 1 minute it indicated that the extract contained saponin (**Adegoke *et al.*, 2010**).

### **Preliminary Trial**

Seeds of soybean var. Anjasmoro were planted in black polythene bags containing 8 kg of alluvial soil and placed in an area surrounded with screen cloth walls and a netting roof for insect protection. 7 days after sprouting the seedlings were fertilized with 0.5g Urea, 1.2g TSP dan 1.2g KCl per bag. Dimethyl sulphoxide was added to crude extract to make it soluble in water. Three concentration levels (25 mg/l, 50 mg/l, 100 mg/l) and a plain water control was then sprayed onto the leaves 14 days after planting at a rate of 25 ml per plant. Each solution was sprayed onto three plants (**Kalaivanan *et al.*, 2012**; **Grabowska *et al.*, 2012**).

### **The Second Trial**

The crude extracts that proved most promising in terms of growth enhancement from the preliminary trial were then trialed again. *C.asiatica*, cassava and *G.linearis* leaf extracts were trailed at four concentration levels (25 mg/l, 50 mg/l, 100 mg/l and 200 mg/l) along with a plain water control. Nested design was used with four replications.



## Data Analysis

The parameters of vegetative growth, plant height, number of leaves, leaf area, dry weight of crown and root and total dry weight of plant, of the second trial were analyzed using ANOVA and Duncan's Multiple Range Test ( $p \leq 0.05$ ).

## RESULTS AND DISCUSSION

The results of phytochemical analysis are shown in Table 1. Each extract contained at least three different classes of secondary metabolites that could have a potential impact on soybean plant growth. It was decided that all six plant extracts warranted further investigation. **Ismail *et al.*, (2016)** reported that both the methanolic and aqueous extracts of *Faidherbia albida* legumes contain terpenes, while alkaloids, saponins and phenolic compounds were found in the aqueous extract, and steroids were found in methanolic extracts only.

Table 1. Results of phytochemical analysis to contents of secondary metabolites in crude extract of six plants

| The source of crude extract      | Flavonoid | Steroid | Terpenoid | Alkaloid | Polar phenolic compound | Semipolar phenolic compound | Saponin |
|----------------------------------|-----------|---------|-----------|----------|-------------------------|-----------------------------|---------|
| Cassava leaf                     | +         | -       | +         | -        | +                       | +                           | -       |
| <i>G. linearis</i> leaf          | +         | -       | +         | -        | -                       | +                           | +       |
| <i>C. asiatica</i>               | -         | +       | +         | -        | +                       | -                           | -       |
| Stigma of corn female flower     | +         | +       | -         | -        | +                       | +                           | -       |
| Bark of <i>A. scholaris</i> stem | -         | +       | +         | +        | +                       | +                           | +       |
| Pericarp of mangosteen fruit     | -         | -       | +         | -        | +                       | +                           | +       |

Note : (+) = present, (-) = absent

Results of the preliminary trial showed, just as **Talukder *et al.*, (2015)** reported, that some plant crude extracts inhibited and others promoted growth. Figures 1 and Figure 2 (a) and 2 (b) shows that the treatment with crude extracts of *C. asiatica*, cassava and *G. linearis* leaves with low concentration showed a tendency to increase plant height and leaf area at 28



days after planting, while crude extracts of stem bark of *A.scholaris* and pericarp of mangosteen fruits and corn silk showed a tendency to decrease the growth parameters.




Figure 1. Plant height of soybean by crude extract treatment of six plants with four concentration levels at 28 days after planting.




Figure 2. Number (a) and Leaf area (b) of soybean by crude extract treatment of six plants with four concentration levels at 28 days after planting.

Notes: CL = cassava leaf; GL = *G. linearis* leaf; CA = *C. asiatica*; CS = stigma of corn female flower; BA = bark of *A.scholaris* stem; PM = pericarp of mangosteen fruit; 1 = control; 2 = 25 mg/l; 3 = 50 mg/l; and 4 = 100 mg/l

A tendency of a crude extract to increase and decrease vegetative growth appears to be related to the secondary metabolites it contains. The crude extracts of *C.asiatica*, cassava and *G.linearis* leaves, that showed the increase of vegetative growth, all contained terpenoids. It is known that at least gibberellin phytohormones, a common class of



diterpenoid, has growth-promoting bioactivity and other terpenoids can promote the action of gibberellins (**Zi *et al.*, 2014**).

The crude extract of *A. scholaris* stem bark and pericarp of mangosteen fruits decreased vegetative growth. Alkaloid and both polar and semipolar phenolic compounds are known to be inhibitors of plant growth. No sign of alkaloids was found in the three extracts that appeared to be growth promoting while all three extracts that appeared to be growth inhibiting contained detectable polar and semipolar phenolic compounds. In general, alkaloid compounds have an allelopathic effect that inhibits growth of both monocotyl and dicotyl plants (**Agbagwa, 2014**).

Polar phenolic compounds have a phytotoxic effect and inhibit cell division (**Zhao-Hui *et al.*, 2010**). While each of the three growth promoting extracts contained either polar or semi-polar phenolic compounds, it is possible that these compounds were present in small enough concentrations that these effects were not as apparent as in the other extracts or were modified by the presence of other metabolites.

Saponin is present in some, but not all of the growth enhancing extracts but it is also present in some of the growth inhibiting ones also. Sometimes saponin has a toxic effect on plants and it inhibits seedling growth (**Prokofeva *et al.*, 2003**; **Kohata *et al.*, 2004**), but saponin triterpenoid shows a growth stimulant activity on corn (**Saha *et al.*, 2010**).

Some concentrations of the crude extracts of the three plants that appeared to promote growth (*C. asiatica*, cassava leaves and leaves of *G. linearis*) were tested again on new soybean plants. The results indicated that some concentrations of these extracts had a statistically significant influence on growth as shown in Table 2.

The only treatment that had a significant effect on leaf area was 25 mg/l crude extract of *C. asiatica*. Other studies have discovered leaf area increases with use of crude extracts. **Agbagwa *et al.*, (2003)** reported seedlings of *Celosia argentea* sprayed with the higher concentrations of the crude extract (75% and 100%) of *Senna alata* showed significant promotion effects on leaflet enlargement. **Fathima *et al.*, (2011)** reported that application of the crude brassinosteroid extract resulted in increase of shoot length, fresh weight, dry weight along with leaf area of seedlings of *Gossypium hirsutum* and *Vigna mungo*. Agbagwa





reported that the growth promoting substances in a crude extract of *Senna alata* caused a general increase in leaf area of *Celosia argentea*L. as well as its height (Agbagwa, 2014).

Table 2. Vegetative growth of soybean resulted by crude extract treatment of cassava leaf, *C. asiatica* and *G. linearis* leaf with five concentration levels at 28 days after planting

| Treatment                   |                                       | Average of           |                    |                              |                         |                         |                               |
|-----------------------------|---------------------------------------|----------------------|--------------------|------------------------------|-------------------------|-------------------------|-------------------------------|
| The source of crude extract | Concentration of crude extract (mg/l) | Plant height(cm)     | Number of leaf     | Leaf area (cm <sup>2</sup> ) | Dry weight of crown (g) | Dry weight of roots (g) | Total dry weight of plant (g) |
| Cassava leaf                | 0 (control)                           | 0.3563 <sup>ab</sup> | 7.00 <sup>ns</sup> | 22.93 <sup>a</sup>           | 1.88 <sup>ns</sup>      | 0.51 <sup>ns</sup>      | 2.40 <sup>ns</sup>            |
|                             | 25                                    | 0.3700 <sup>c</sup>  | 7.00 <sup>ns</sup> | 26.51 <sup>a</sup>           | 2.08 <sup>ns</sup>      | 0.54 <sup>ns</sup>      | 2.62 <sup>ns</sup>            |
|                             | 50                                    | 0.3700 <sup>c</sup>  | 7.25 <sup>ns</sup> | 27.31 <sup>a</sup>           | 2.23 <sup>ns</sup>      | 0.53 <sup>ns</sup>      | 2.76 <sup>ns</sup>            |
|                             | 100                                   | 0.3600 <sup>b</sup>  | 7.00 <sup>ns</sup> | 23.50 <sup>a</sup>           | 2.04 <sup>ns</sup>      | 0.50 <sup>ns</sup>      | 2.55 <sup>ns</sup>            |
|                             | 200                                   | 0.3550 <sup>a</sup>  | 6.75 <sup>ns</sup> | 22.55 <sup>a</sup>           | 1.93 <sup>ns</sup>      | 0.54 <sup>ns</sup>      | 2.47 <sup>ns</sup>            |
| <i>C. asiatica</i>          | 0 (control)                           | 0.3725 <sup>c</sup>  | 7.00 <sup>ns</sup> | 26.53 <sup>a</sup>           | 2.08 <sup>ns</sup>      | 0.53 <sup>ns</sup>      | 2.61 <sup>ns</sup>            |
|                             | 25                                    | 0.4573 <sup>d</sup>  | 7.25 <sup>ns</sup> | 33.01 <sup>b</sup>           | 2.46 <sup>ns</sup>      | 0.61 <sup>ns</sup>      | 3.07 <sup>ns</sup>            |
|                             | 50                                    | 0.3575 <sup>b</sup>  | 7.00 <sup>ns</sup> | 27.33 <sup>a</sup>           | 2.03 <sup>ns</sup>      | 0.50 <sup>ns</sup>      | 2.53 <sup>ns</sup>            |
|                             | 100                                   | 0.3538 <sup>a</sup>  | 6.75 <sup>ns</sup> | 24.01 <sup>a</sup>           | 1.75 <sup>ns</sup>      | 0.36 <sup>ns</sup>      | 2.11 <sup>ns</sup>            |
|                             | 200                                   | 0.3738 <sup>c</sup>  | 6.75 <sup>ns</sup> | 25.59 <sup>a</sup>           | 1.58 <sup>ns</sup>      | 0.56 <sup>ns</sup>      | 2.15 <sup>ns</sup>            |
| <i>G. linearis</i> leaf     | 0 (control)                           | 0.3588 <sup>bc</sup> | 6.75 <sup>ns</sup> | 25.38 <sup>a</sup>           | 1.87 <sup>ns</sup>      | 0.56 <sup>ns</sup>      | 2.43 <sup>ns</sup>            |
|                             | 25                                    | 0.3425 <sup>a</sup>  | 6.50 <sup>ns</sup> | 25.87 <sup>a</sup>           | 2.03 <sup>ns</sup>      | 0.50 <sup>ns</sup>      | 2.53 <sup>ns</sup>            |
|                             | 50                                    | 0.3563 <sup>b</sup>  | 7.00 <sup>ns</sup> | 26.28 <sup>a</sup>           | 2.06 <sup>ns</sup>      | 0.54 <sup>ns</sup>      | 2.60 <sup>ns</sup>            |
|                             | 100                                   | 0.3638 <sup>d</sup>  | 6.75 <sup>ns</sup> | 27.72 <sup>a</sup>           | 1.92 <sup>ns</sup>      | 0.56 <sup>ns</sup>      | 2.48 <sup>ns</sup>            |
|                             | 200                                   | 0.3563 <sup>b</sup>  | 7.25 <sup>ns</sup> | 22.86 <sup>a</sup>           | 1.75 <sup>ns</sup>      | 0.57 <sup>ns</sup>      | 2.31 <sup>ns</sup>            |

Note: values with common letter in the same column are not significantly different at 5% level as per DMRT; ns = non significant

Treatment using 25 mg/l crude extract of *C. asiatica* resulted in significantly higher plant height than the control and the other concentrations of *C. asiatica*. This extract increase in plant height was observed between 14 to 28 days after planting, while the plant height at other concentrations was relatively similar to the control treatment at every week as can be seen from Fig. 4. Table 2 also shows that use of 25 mg/l and 50 mg/l concentrations of cassava leaf and 100 mg/l of *G. linearis* leaf extracts also produced higher plants although the growth enhancement compared to the control was not as great as with *C. asiatica*.






Figure 4. Plant height of soybean resulted treatment of *C. asiatica* crude extract with five concentration levels from 7 up to 28 day after planting. Note: dap = days after planting

Interestingly, *C. asiatica* crude extract with low concentration (25 mg/l) increased plant height and leaf area while high concentration decreased these parameters. The concentration of crude extracts of plant material has been shown to be an influential factor in studies using much higher concentrations. **Emongor (2014)** reported that *Moringa* leaf extract at 11% up to 50% concentration increased vegetative growth of snap beans (*Phaseolus vulgaris*) but concentrations outside this range did not. **Baeshen (2014)** reported that aqueous extract of *Mentha peperina* which could enhance the seed germination and other growth parameters of *Lens culinary* with a lower concentration of crude extracts, but these were reduced at a higher one.

## CONCLUSIONS

The results showed that low concentration crude extract of *C. asiatica* (25 mg/l) significantly increased both height and leaf area of soybean plants. The height was 22.77% higher than the control. Low concentration of crude extract is a potential growth stimulant for soybean plants. Further investigations will determine the effect of steroids and terpenoids from purified extracts of *C. asiatica* on the growth and production of soybean. If these secondary metabolites are found to be effective growth stimulants, this will help in the search for an environmentally friendly, commercially viable and cost effective way to improve soybean production in a tropical context.



## ACKNOWLEDGEMENTS

The financial support from Andalas University is gratefully appreciated. The author also acknowledges rector for providing the professor research grant which involved a student of the doctoral program.

## REFERENCES

- [1]. Abdalla, M.M. 2014. Boosting the growth of rocket (*Eruca vesicaria* subsp. *sativa*) plants in response to the application of *Moringa oleifera* extracts as biostimulant. Life Sci.J.11(11): 1113-1121. <http://lifesciencesite.com.189>
- [2]. Adegoke, A.A., P.A. Iberi, D.A. Akinpelu, O.A. Aiyegoro and C.I. Mbotto. 2010. Studies on phytochemical screening and antimicrobial potentials of *Phyllanthus amarus* against multiple antibiotic resistant bacteria. Int. J. Appl. Res. Nat. Prod. 3 (3): 6–12.
- [3]. Agbagwa, I.O, F.A. Onofeghara and S.I. Mensah. 2003. Stimulation of growth and development of *Celosia argentea* L. by crude extracts of *Senna alata* (L.) Roxb. J. Appl. Sci. Environ. Mgt. 7(1): 9-13.
- [4]. Agbagwa, I.O. 2014. Crude extracts of *Senna alata*(L.) Roxb.mimics plant growth hormones in promotion of vegetative and reproductive growth in *Celosia argentea* L. AM J. Plant Sci., 5: 1918-1925.<http://dx.doi.org/10.4236/ajps.2014.513205>
- [5]. Anbuselvi, A. and T. Balamurugan. 2014. Phytochemical and antinutrient constituents of cassava and sweet potato. WJPPS 3(3): 14-40-1449.
- [6]. Baeshen, A.A. 2014. Assessment of the effect of some medicinal plants extracts on germination and growth of lentil (*Lens culinaris*). Re. J. Pharm. Biol. Chem. Sci., 5(3): 1846-1853.
- [7]. Bhaigyabati, T., T. Kirithika, J. Ramya and K. Usha. 2011. Phytochemical constituents and antioxidant activity of various extracts of corn silk (*Zea mays* L.). Res. J. Pharm. Biol. Chem. Sci. 2(4): 986-993.
- [8]. Biradar, S.R., and B.D. Rachetti. 2013. Extraction of some secondary metabolites and thinlayer chromatography from different parts of *Centella asiatica* L.(URB). American J.Life Sci. 1(6):243-247 . doi: 10.11648/j.ajls.20130106.11
- [9]. Boxi, M., Y. Rajesh, V.R. Kumar, B. Praveen and K. Mangamma. 2010. Extraction, phytochemical screening and in-vitro evaluation of anti-oxidant properties of *Commicarpus schinensis* (aqueous leaf extract). Int. J. Pharm. Biosci. 1: 537–547.
- [10]. Culver, M., T. Fanuel and A.Z. Chiteka. 2012. Effect of Moringa extract on growth and yield of tomato. Greener J. Agric. Sci. 2(5): 207-211.
- [11]. Demissie, A.G. and S.S. Lele. 2013. Bioactivity-directed isolation and identification of novel alkaloid from *Jatropha curcas* (Linn.). RJCES 1(2): 22-28.
- [12]. Dey, A. 2011.*Alstoniascholaris* R.Br. (Apocynaceae): Phytochemistry and pharmacology: A concise review. J. Appl. Pharm. Sci. 1(6): 51-57.
- [13]. Directorate General of Crops. 2014. Program and Activity of Crops Development in 2015-2019.The Meeting of Development Plant of Agriculture in 2015.Jakarta, Mei 13<sup>th</sup> 2014. Ministry of Agriculture, Republic of Indonesia.



- [14]. Emongor, V.E. 2014. Effects of *Moringa (Moringa oleifera)* leaf extract on growth, yield and yield components of snap beans (*Phaseolus vulgaris*). *British J. Appl. Sci. Tech.* 6(2): 114-122.
- [15]. Ertani,A., P.Samlo, C.Nicoletto, S.Santagata, M.Schiavon and S.Nardi. 2015. The use of organic biostimulants in hot pepper plants to help low input sustainable agriculture. *Chem.and Biol.Tech.in Agr.* 2(11) :1-10. doi : 10.1186/s40538-015-0039-z.
- [16]. Ertani, A., D. Pizzeeghello, O. Francioso, A. Tinti and S. Nardi. 2016. Biological Activity of Vegetal Extracts Containing Phenols on Plant Metabolism. *Molecules*, 250. DOI: 103390/molecules21020205. [www.mdpi.com/journal/molecules](http://www.mdpi.com/journal/molecules).
- [17]. Fathima, S.A., M. Johnson and K. Lingakumar. 2011. Effect of crude brassinosteroid extract on growth and biochemical changes of *Gossypiumhirsutum*L. And *Vignamungo*L. *J. Stress Physiol.Biochem.* 7(4): 324-334.
- [18]. Grabowska, A., E. Kunicki, A. Sekara and A.Kalisz. 2012. The effect of cultivar and biostimulant treatment on the carrot yield and its quality. *Veg. Crops Re. Bull.* 77: 37-48.
- [19]. Gutierrez-Orozco, F. and M.L. Failla. 2013. Biological activities and bioavailability of mangosteenxanthones. *Nutrients*, 5(8): 3163-3183.
- [20]. Hanafy, M.S., F.M. Saadawy, S.M.N. Milad and R.M. Ali. 2012. Effect of some natural extracts on growth and chemical constituents of *Scheffleraarboricolap*plants. *J. Hort. Sci.Orn. Plants* 4(1): 26-33.
- [21]. Hasanudin, K., P.Hashim and S.Mustafa. 2012. Corn Silk (stigma maydis) in Healthcare : A phytochemical and Pharmacological Review. 2012. *Molecules*. 17 : 9697-9715. Doi: 10.3390/molecules 17089697
- [22]. Ismail, A.M., E.A. Mohamed, M.R. Marghany, F.F. Abdel-Motaal, I.B. Abdel-Farid, M.A. El-Sayed. 2016. Preliminary phytochemical screening, plant growth inhibition and antimicrobial activity studies of *Faidherbiaalbida* legume extracts. *J. Saudi Soc. Agric. Sci.*15(2): 112–117.
- [23]. James, J.T. and I.A. Dubery. 2011. Identification and quantification of triterpenoidcentelloids in *Centellaasiatica* (L.) Urban by densitometric TLC. *J. Planar Chromatog.* 24(1): 82-87.
- [24]. Jubahar, J., Dachriyanus, D. Arbain, A. Bakhtiar, H. Mukhtar, and M.V. Sargent. 2006. A Flavonoid sulphate from *Gleichenialinear*is (Burm; Clarke). *Asian Coordinating Group for Chemistry (ACGC) Chemical Research Communication* 20: 6-7.
- [25]. Kalaivanan, C., M. Chandrasekaran, and V. Venkatesalu. 2012. Effect of seaweed liquid extract of *Caulerpascalpelliformis* on growth and biochemical constituents of black gram (*Vigna radiate* L.(Hepper). *Phykos*, 42(2): 46-53.
- [26]. Kohata, K., Y. Yamauchi, T. Ujihara and H. Horie. 2004. Growth inhibitory of tea-seed saponis and glyphosate to weed seedling. *JARQ* 38(4): 267-270.
- [27]. Muhamman, M.A. and S.G.G. Mohammed. 2014. Effects of aqueous extract of moringa (*Moringaoleifera*L.) and nitrogen rates on the contribution of some physiological and yield attributes to grain yield of sesame (*Sesamumindicum*L.) II. *Sci. Papers. Series A. Agronomy*, Vol. LVII: 272-275.
- [28]. Mvumi, C., F. Tagwira and A.Z. Chiteka. 2012. Effect of moringa extract on growth and yield of maize and common beans. *Greener J. Agric. Sci.* 3(1): 055-062.
- [29]. Nuwamanya, E., P.R. Rubaihayo, S. Mukasa, S. Kyamanyawa, J.F. Huwamba and Y. Baguma. 2014. Biochemical and secondary metabolites changes under moisture and temperature stress in cassava (*Manihotesculenta*Crantz). *African.J.of Biotech.* 13(31): 3173-3186.



- [30]. Obianime, A.W. and F.I. Uche. 2008. The phytochemical screening and effects of methanolic extract of *Phyllanthusamarus* leaf on the biochemical parameters of male guinea pigs. J. Appl. Sci. Environ. Manage. 12 (4): 73–77.
- [31]. Pankti, K., G. Payal, C. Manodeep and K. Jagadish. 2012. A phytopharmacological review of *Alstoniascholaris*: A Panoramic herbal medicine. Intl. J. Re.Ayuveda and Pharm.(3(3): 367-371.
- [32]. Pratap, B., G.S. Chakraborty and Mogha. 2013. Complete aspects of *Alstoniascholaris*. Intl. J.Pharm.Tech. Re. 5(1): 17-26.
- [33]. Prokofeva, NG., E.L. Chaikina, A.A. Kicha, N.V. Ivanchina. 2003. Biological activities of steroid glycosides from starfish.Compr.Biochem. Physiol. B: Biochem. Mol. Biol. 137: 695-701.
- [34]. Saha, S., S. Walia, J. Kumar and B.S. Parmar. 2010. Triterpenic saponins as regulator of plant growth. J. Apl. Bot. Food Qlty. 83: 189-195.
- [35]. Sarmin, N.S., 2014. Effect of *Moringaoleiferaon* germination and growth of *Triticumaestivum*.J.Biosci. Agric. Re.02(02): 59-69
- [36]. Shahid-Ud-Duaula, A.F.M. and B.M. Anwarul. 2009. Phytochemical screening, plant growth inhibition, and antimicrobial activity studies of *Xylocarpusgranatum*.Malaysian. J. Pharm. Sci. 7(1): 9–21.
- [37]. Shao, H., X. Huang, Y. Zhang and C. Zhang. 2013. Main alkaloids of *Peganumharmala* L. and their different effects on dicot and monocot crops. Molecules,18: 2623-2634.
- [38]. Singh, J., P. Singh, A. Gupta, S. Solanki, E. Sharmaand, R.Nema. 2012. Qualitative estimation of the presence of bioactive compound in *Centellaasiatica*: An important medicinal plant. Intl. J. Life Sci. Med. Sci. 2(1): 5-7.
- [39]. Siyumbawe, S., N. Nomishan, Beny and P.N. Okechukwu. 2014. Antioxidant and gastro-protective activity of ethanolic rind extract of *Manihotesculenta*Crantz. Intl. J. Pharm. Re.Schlr. 3(1-2):798-808.
- [40]. Talukder, M.A.I., M. Rahaman, B. Roy and K.C. Saha. 2015. Effects of herbal plantextracts on germination and seedling growth of some vegetables. Intl. J. Sci. Nat. 6 (3): 421-425.
- [41]. Vriet, C., E. Russinova and C. Reuzeau. 2012. Boosting crop yields with plant steroids. Plant Cell 24(3): 842–857.
- [42]. Yusuf, U., F. Abdullah, K. Itam and M.A. Sukari. 2003. Leaf flavonoids in the genus *Gleichenia* (Gleicheniaceae). Am. Fern J. 93(1): 44-46.
- [43]. Zarena, A.S. and K.U. Sakar. 2012. Phenolic acid, flavonoid profile and antioxidant activity in mangosteen (*Garciniamangostana*L.) pericarp. J. Food Biochem. 36(5): 627-633.
- [44]. Zhao-Hui, L., W. Qiang, R. Xiao, P. Cun-De and J. De-An. 2010. Phenolics and plant allelopathy. Molecules 15: 8933-8952.
- [45]. Zi, J., S.Mafu and R.J. Peters.2014. To Gibberellins and beyond! surveying the evolution of (di)terpenoid metabolism. Annl.Rev. Plant Biol. 65: 259–286.