



# Plant Geometry, Macro and Micro Nutrients on Growth and Growth Analysis of Dual Purpose Sorghum under Rainfed Vertisol Condition

G. Bharathi<sup>1\*</sup>, M. Joseph<sup>2</sup>, A. Velayutham<sup>3</sup>, K. Baskar<sup>4</sup>

<sup>1</sup>\*PG Scholar

<sup>2</sup>Assistant Professor, Department of Agronomy

<sup>3</sup>Professor, Department of Agronomy

Agricultural College and Research Institute, Killikulam, Thoothukudi District, Tamil Nadu

<sup>4</sup>Professor, Department of Soil Science and Agricultural Chemistry, ARS, Kovilpatti

[bharathiagri04@gmail.com](mailto:bharathiagri04@gmail.com); 7373592613

\*Corresponding Author

**Abstract:** Field experiment was conducted at Agricultural Research Station, Kovilpatti during rabi season (October 2017 to January 2018) to find out the suitable plant geometry, levels of fertilizers and foliar spray of micronutrients (Zn and Fe) at 15, 30 and 45 DAS to improve the growth and yield of sorghum variety K12 (dual purpose). The experiment was laid out in randomized block design with three replication and twelve treatments. The treatment consists of two plant geometry, three different dose of fertilizer and foliar spray of micronutrients viz., 0.5% ZnSO<sub>4</sub> + 0.2% FeSO<sub>4</sub> at 15, 30 and 45 DAS. Observation on growth parameters, growth analysis and finally grain and stover yield were recorded. The experimental results shows that reduced plant spacing of 30 × 15 cm together with enhanced application of 50:25:25 kg NPK ha<sup>-1</sup> + foliar spray of 0.5% ZnSO<sub>4</sub> + 0.2% FeSO<sub>4</sub> at 15, 30 and 45 DAS significantly increased the growth parameters viz., plant height, stem girth and days to flowering. This attributes ultimately resulted in higher grain (3961 kg ha<sup>-1</sup>) and stover (13972 kg ha<sup>-1</sup>) yield of dual purpose sorghum compared to recommended practice of rainfed sorghum having 45 × 15 cm spacing and application of recommended 40:20:0 kg NPK ha<sup>-1</sup> alone.

**Keywords:** Plant spacing - Soil & Foliar nutrition – Growth and yield - Rainfed sorghum



## INTRODUCTION

Sorghum [*Sorghum bicolor* (L.) Moench] is one of the gifted grass genera of tropics and in India, it is popularly known as “Jowar”. Its grains have about 10-12% protein, 3% fat and 70% carbohydrate. It provides food, feed and fuel to millions of poor farm families and their livestock in the arid and semi-arid tropical region of the world. India is one of the notable country for its higher livestock population and its economic integration with farm production, especially under the less mechanized dry land agriculture. Sorghum is most important crop to resource poor farmers for nutritional and livelihood security. The demand and consumption of agricultural crops for food, feed, and fuel is increasing at a globally rapid pace. Therefore, the production practices must be well optimized and managed to satisfy the growing worldwide demand. In more advanced agriculture the grain is used as a livestock feed while the whole crop is used as silage or forage (Cobley, 1976). The growth and yield of a crop can be adversely affected by deficient or excessive supply of any one of the essential nutrients. Soil application of nutrients will give the initial boost for growing seedlings (Hamayun *et al.*, 2011). Growth components like plant height, number of leaves, stem diameter, leaf area of fodder maize influenced significantly by the application of nitrogen and phosphorus (Ayub *et al.*, 2002). Hence this study was initiated with the objectives of determining the effect of different plant geometry, fertilizers levels and foliar spray of micronutrients on the performance of dual purpose sorghum under rainfed conditions.



## Materials and Methods

The field experiment was conducted at Agricultural Research station, Kovilpatti during *rabi* (Oct 2017 – Jan 2018). The soil was clay loam in texture with pH of 8.04, EC of 0.45 dS m<sup>-1</sup> and organic carbon content of 3.38 g kg<sup>-1</sup>. The soil was low in available nitrogen (179 kg ha<sup>-1</sup>), low in available phosphorus (10 kg ha<sup>-1</sup>) and high in available potassium (365 kg ha<sup>-1</sup>). The mean annual rainfall is 703 mm and the maximum and minimum temperature ranges from 34.9°C and 22.8°C, respectively. During the cropping period, the crop received 421.5 mm of rainfall in 17 rainy days and the maximum and minimum temperature ranges from 36.5°C and 17.6°C, respectively.

The experiment was laid out in randomized block design with three replication and twelve treatments. The treatment consists of two plant geometry, three different dose of fertilizer and foliar spray of micronutrients *viz.*, 0.5% ZnSO<sub>4</sub> + 0.2% FeSO<sub>4</sub> at 15, 30 and 45 DAS. The sorghum variety K 12 (dual purpose) was chosen for this study. All agronomic practices like weed control, thinning, plant protection measures and harvesting as well as postharvest operation were made uniformly for all treatments. Various observation *viz.*, plant height, stem girth, days to flowering, earhead length, earhead breadth, number of filled grains earhead<sup>-1</sup>, test weight, finally grain and stover yield were recorded.



## Result and Discussion

### Effect of treatments on growth parameters

Growth parameters of rainfed dual purpose sorghum was significantly influenced by adoption of different treatment combinations (Table 1). The plant height of various treatments ranged from 201.5 cm to 238.9 cm. The maximum plant height of 238.9 cm at harvest was recorded in the plant spacing of  $30 \times 15$  cm with basal application of 50:25:25 kg NPK ha<sup>-1</sup> + foliar spray of 0.5% ZnSO<sub>4</sub> + 0.2% FeSO<sub>4</sub> at 15, 30 and 45 DAS. Competition for light might be the responsible for increase in height due to closer intra-row spacing in sorghum. This statement was agrees with the results of Miko and Manga, 2008. Increased nitrogen application also increased the plant height might be due to the progressive effect of nitrogen element on plant growth that leads to progressive increase in inter nodal length and consequently plant height. Similar results were reported by Gupta *et al.* (2008) and Nirmal *et al.* (2016).

Among different spacing,  $45 \times 15$  cm (T<sub>1</sub>-T<sub>6</sub>) recorded higher mean stem girth value of 2.48 cm than the closer spacing of  $30 \times 15$  cm (T<sub>7</sub>-T<sub>12</sub>) recorded the mean of 2.28 cm. The stem girth was higher in  $45 \times 15$  cm spacing along with basal application of 50:25:25 kg NPK ha<sup>-1</sup> + foliar spray of 0.5% ZnSO<sub>4</sub> + 0.2% FeSO<sub>4</sub> at 15, 30 and 45 DAS which recorded 2.7 cm. the reason for stem diameter reduction at higher plant densities can be linked to a reduction of assimilates allocation and more intra-plant competition among plants. (Zand and Shakiba, 2013).

Earlier advancement of flowering was observed in the graded levels of NPK application under both the plant populations. Besides, foliar application of micronutrients (0.5% ZnSO<sub>4</sub> +



0.2% FeSO<sub>4</sub> at 15, 30 and 45 DAS) was found to delay the flowering in dual purpose rainfed sorghum than the basal application of NPK alone in irrespective of the sowing geometry.

Sorghum K12 was early flowering type variety. Usually flowering arises 60 days after sowing. Higher level of nutrients and foliar spray of micronutrients in reduced plant populations delayed the flowering. This could be related to the supportive effects of more available fertilizers to lower number of plants unit area<sup>-1</sup> which permitted building of more vigorous growth that resulted in higher number of days to flowering. More plant density resulted inter competition increased and early flowering. These results were close conformity with Zand and Shakiba, (2013).

Dry matter production is basically a measure of photosynthetic efficiency of assimilatory system in plants. Plant spacing of 30 × 15 cm coupled with graded level of NPK @ 50:25:25 kg ha<sup>-1</sup> + foliar spray of 0.5% ZnSO<sub>4</sub> + 0.2% FeSO<sub>4</sub> at 15, 30 and 45 DAS was found to accumulate more drymatter of 18254 kg ha<sup>-1</sup> at harvest stage (Table 2). This might be due to increase in nitrogen level having resulted in more active plant growth, which consecutively resulted in more drymatter partitioning. Similar results were also reported by Singh *et al.* (2005).

### **Effect of treatments on growth analysis**

Plants with more leaves produced higher leaf area index. Maximum leaf area index was noticed in 45 × 15 cm spacing with basal application of 50:25:25 kg NPK ha<sup>-1</sup> + foliar spray of 0.5% ZnSO<sub>4</sub> + 0.2% FeSO<sub>4</sub> at 15, 30 and 45 DAS recorded 5.94 at 30 DAS and 11.85 at 60



DAS (Table 3). The significant increase in LAI possibly due to the improved leaf number and expansion in plants and application of nitrogenous fertilizers (Shamme *et al.*, 2016).

The crop growth rate of sorghum ranged from 154.26 to 167.99 kg ha<sup>-1</sup> day<sup>-1</sup> under 45 × 15 cm plant spacing and from 159.18 to 192.96 kg ha<sup>-1</sup> day<sup>-1</sup> under 30 × 15 cm plant spacing at harvest stage (Table 4). Crop growth rate showed a rapid increase from sowing to 60 DAS and thereafter slow increase up to harvest. Plant spacing of 30 × 15 cm with graded level of NPK @ 50:25:25 kg ha<sup>-1</sup> + foliar spray of 0.5% ZnSO<sub>4</sub> + 0.2% FeSO<sub>4</sub> at 15, 30 and 45 DAS was recorded higher CGR of 66.23, 317.12 and 192.96 kg ha<sup>-1</sup> day<sup>-1</sup> at 30 DAS, 60 DAS and at harvest stages, respectively. The treatments with graded levels of NPK alone performed equal and higher growth rate than the treatments having recommended NPK coupled with foliar nutrition.

### **Effect of treatments on yield**

The yield of rainfed dual sorghum K12 was significantly influenced by different plant geometry, graded levels of fertilizer and micronutrient foliar application (Table 5). Grain yield depends on the synthesis and accumulation of photosynthates and their distribution among various plant parts. The synthesis, accumulation and translocation of photosynthates depend upon efficient photosynthetic structure as well as the extent of translocation into sink (grains) and also on plant growth and development during early stages of crop growth. A distinct enhancement in straw yield with increase in nutrient levels was evident of this study. Stover yield is directly proportional to plant population and drymatter accumulation in K12 sorghum.



Sowing at a plant spacing of  $30 \times 15$  cm coupled with graded fertilizer level of 50:25:25 kg NPK  $\text{ha}^{-1}$  + foliar spray of 0.5%  $\text{ZnSO}_4$  + 0.2%  $\text{FeSO}_4$  at 15, 30 and 45 DAS recorded significantly higher grain yield ( $3961 \text{ kg ha}^{-1}$ ) of rainfed sorghum. The increase in grain yield could be as a result of good drymatter production for grain filling as a result of higher number of leaves. Increase in yield of sorghum also due to nitrogen application and has been reported by some workers, Obilana (1983) and Galbiatti *et al.* (1977). The recommended practices for rainfed sorghum having  $45 \times 15$  cm spacing with basal application of 40:20:0 kg NPK  $\text{ha}^{-1}$  recorded 35.0 per cent reduced grain yield compared to the above said best treatment.

Stover yield of rainfed sorghum was also significantly influenced by plant spacing, nutrient levels and foliar spray. Sowing at  $30 \times 15$  cm with higher levels of NPK @ 50:25:25 kg  $\text{ha}^{-1}$  + foliar spray of 0.5%  $\text{ZnSO}_4$  + 0.2%  $\text{FeSO}_4$  at 15, 30 and 45 DAS recorded significantly higher stover yield of  $13972 \text{ kg ha}^{-1}$  (Table 5). The significant increase in fodder yield with increasing fertility levels was due to fact that all these nutrients were involved in increasing protoplasmic constituents, root, shoot growth and accelerating the process of cell division, enlargement and elongation which in turn showed luxuriant vegetative growth and resulted in higher green and dry fodder yield. Similar results were also obtained by Kumar and Chaplot (2015) and Rana *et al.*(2013).



## REFERENCES

- [1]. Ayub, M., Tanveer, A., Ali, S., & Nadeem, M. A. (2002). Effect of different nitrogen levels and seed rates on growth, yield and quality of sorghum (*Sorghum bicolor*) fodder. *Indian journal of agricultural science*, 72(11), 648-650.
- [2]. Cobby, L. S. (1976). *An introduction to the botany of tropical crops*: Longman Group Limited.
- [3]. Galbiatti, J. A., Benincasa, M. M. P. And Benincas, M. (1977). Effect of rates and date of ammonium sulphate application on the prefer income of *Sorghum bicolor* L. Moench. *Field Crops Abstracts*, 1246.
- [4]. Gupta, K., Rana, D., & Sheoran, R. (2008). Response of nitrogen and phosphorus levels on forage yield and quality of sorghum [*Sorghum bicolor* (L.) Moench]. *Forage Res*, 34(3), 156-159.
- [5]. Hamayun, M., Khan, S. A., Khan, A. L., Shinwari, Z. K., Ahmad, N., Kim, Y.-H., & Lee, I.-J. (2011). Effect of foliar and soil application of nitrogen, phosphorus and potassium on yield components of lentil. *Pak. J. Bot*, 43(1), 391-396.
- [6]. Kumar, D., & Chaplot, P. (2015). Performance of multicut forage sorghum genotypes to fertility levels. *Forage Res*, 41, 199-201.
- [7]. Miko, S., & Manga, A. (2008). Effect of intra-spacing and nitrogen rates on growth and yield of sorghum (*Sorghum bicolor* L.) var. ICSV 400. *Prod. Agric. Technol. J*, 4, 66-73.





- [8]. Nirmal, S., Dudhade, D., Solanke, A., Gadakh, S., Bhakare, B., Hasure, R., & Gore, S. (2016). Effect of nitrogen levels on growth and yield of forage sorghum [*Sorghum bicolor* (L.) Moench] varieties. *International Journal of Science, Environment and Technology*, 5(5), 2999-3004.
- [9]. Obilana, A. T. (1983). *Status of sorghum research towards better increase productivity*. Paper presented at the First National WorkShop of NAFPP, Ibadan.
- [10]. Rana, D., Singh, B., Gupta, K., Dhaka, A., & Pahuja, S. (2013). Effect of fertility levels on growth, yield and quality of multicut forage sorghum [*Sorghum bicolor* (L.) Moench] genotypes. *Forage Res*, 39, 36-38.
- [11]. Shamme, K., Raghavaiah, C. V., Balemi, T., & Hamza, I. (2016). Sorghum (*Sorghum bicolor* L.) Growth, Productivity, Nitrogen Removal, N-Use Efficiencies and Economics in Relation to Genotypes and Nitrogen Nutrition in Kellem-Wollega Zone of Ethiopia, East Africa. *Advances In Crop Science And Technology*.
- [12]. Singh, M., Maurya, M., Singh, S., & Mishra, C. (2005). Effect of nitrogen level and biofertilizer inoculation on productivity of forage sorghum (*Sorghum bicolor*). *Indian journal of agricultural science*, 75(3), 167-168.
- [13]. Zand, N., & Shakiba, M. R. (2013). Effect of plant density and nitrogen fertilizer on some attribute of grain sorghum (*Sorghum bicolor* (L.) moench). *International Journal of Advanced Biological and Biomedical Research*, 1(12), 1577-1582.



**Table 1. Effect of plant geometry, soil and foliar nutrition on plant height (cm), stem girth (cm) and days to flowering of rainfed dual purpose sorghum**

Treatments		plant height (cm) at harvest	Stem girth	Days to flowering	
T <sub>1</sub>	45 × 15 cm spacing	40:20:0 kg NPK ha <sup>-1</sup> (RDF)	201.5	2.2	58.0
T <sub>2</sub>		40:20:0 kg NPK ha <sup>-1</sup> + FS 0.5% ZnSO <sub>4</sub> + 0.2% FeSO <sub>4</sub> at 15, 30 and 45 DAS	208.5	2.3	58.6
T <sub>3</sub>		40:20:20 kg NPK ha <sup>-1</sup>	216.4	2.3	59.0
T <sub>4</sub>		40:20:20 kg NPK ha <sup>-1</sup> + FS 0.5% ZnSO <sub>4</sub> + 0.2% FeSO <sub>4</sub> at 15, 30 and 45 DAS	222.8	2.7	60.0
T <sub>5</sub>		50:25:25 kg NPK ha <sup>-1</sup>	219.2	2.7	60.5
T <sub>6</sub>		50:25:25 kg NPK ha <sup>-1</sup> + FS 0.5% ZnSO <sub>4</sub> + 0.2% FeSO <sub>4</sub> at 15, 30 and 45 DAS	224.3	2.7	62.0
T <sub>7</sub>	30 × 15 cm spacing	40:20:0 kg NPK ha <sup>-1</sup> (RDF)	205.1	2.1	58.0
T <sub>8</sub>		40:20:0 kg NPK ha <sup>-1</sup> + FS 0.5% ZnSO <sub>4</sub> + 0.2% FeSO <sub>4</sub> at 15, 30 and 45 DAS	212.1	2.2	59.0
T <sub>9</sub>		40:20:20 kg NPK ha <sup>-1</sup>	227.5	2.2	59.0
T <sub>10</sub>		40:20:20 kg NPK ha <sup>-1</sup> + FS 0.5% ZnSO <sub>4</sub> + 0.2% FeSO <sub>4</sub> at 15, 30 and 45 DAS	235.2	2.4	59.4
T <sub>11</sub>		50:25:25 kg NPK ha <sup>-1</sup>	231.2	2.4	59.7
T <sub>12</sub>		50:25:25 kg NPK ha <sup>-1</sup> + FS 0.5% ZnSO <sub>4</sub> + 0.2% FeSO <sub>4</sub> at 15, 30 and 45 DAS	238.9	2.4	61.0
SEd		6.72	0.072	1.20	
CD (p=0.05)		14.10	0.15	NS	



**Table 2. Effect of plant geometry, soil and foliar nutrition on drymatter production (kg ha<sup>-1</sup>) of rainfed dual purpose sorghum**

Treatments			Drymatter production (kg ha <sup>-1</sup> )		
			30 DAS	60 DAS	At harvest
T <sub>1</sub>	45 × 15 cm spacing	40:20:0 kg NPK ha <sup>-1</sup> (RDF)	1148	7492	12891
T <sub>2</sub>		40:20:0 kg NPK ha <sup>-1</sup> + FS 0.5% ZnSO <sub>4</sub> + 0.2% FeSO <sub>4</sub> at 15, 30 and 45 DAS	1295	7710	13308
T <sub>3</sub>		40:20:20 kg NPK ha <sup>-1</sup>	1451	8152	13828
T <sub>4</sub>		40:20:20 kg NPK ha <sup>-1</sup> + FS 0.5% ZnSO <sub>4</sub> + 0.2% FeSO <sub>4</sub> at 15, 30 and 45 DAS	1651	9397	15206
T <sub>5</sub>		50:25:25 kg NPK ha <sup>-1</sup>	1564	8806	14489
T <sub>6</sub>		50:25:25 kg NPK ha <sup>-1</sup> + FS 0.5% ZnSO <sub>4</sub> + 0.2% FeSO <sub>4</sub> at 15, 30 and 45 DAS	1687	9691	15571
T <sub>7</sub>	30 × 15 cm Spacing	40:20:0 kg NPK ha <sup>-1</sup> (RDF)	1150	7515	13086
T <sub>8</sub>		40:20:0 kg NPK ha <sup>-1</sup> + FS 0.5% ZnSO <sub>4</sub> + 0.2% FeSO <sub>4</sub> at 15, 30 and 45 DAS	1356	8045	13686
T <sub>9</sub>		40:20:20 kg NPK ha <sup>-1</sup>	1743	9848	15782
T <sub>10</sub>		40:20:20 kg NPK ha <sup>-1</sup> + FS 0.5% ZnSO <sub>4</sub> + 0.2% FeSO <sub>4</sub> at 15, 30 and 45 DAS	1871	10794	17157
T <sub>11</sub>		50:25:25 kg NPK ha <sup>-1</sup>	1841	10583	16849
T <sub>12</sub>		50:25:25 kg NPK ha <sup>-1</sup> + FS 0.5% ZnSO <sub>4</sub> + 0.2% FeSO <sub>4</sub> at 15, 30 and 45 DAS	1987	11501	18254
SEd			45.89	305.04	512.29
CD (p=0.05)			94.17	632.61	1095.78



**Table 3. Effect of plant geometry, soil and foliar nutrition on leaf area index of rainfed dual purpose sorghum**

Treatments			Leaf area index	
			30 DAS	60 DAS
T <sub>1</sub>	45 × 15 cm spacing	40:20:0 kg NPK ha <sup>-1</sup> (RDF)	3.25	8.36
T <sub>2</sub>		40:20:0 kg NPK ha <sup>-1</sup> + FS 0.5% ZnSO <sub>4</sub> + 0.2% FeSO <sub>4</sub> at 15, 30 and 45 DAS	3.87	9.14
T <sub>3</sub>		40:20:20 kg NPK ha <sup>-1</sup>	4.27	9.76
T <sub>4</sub>		40:20:20 kg NPK ha <sup>-1</sup> + FS 0.5% ZnSO <sub>4</sub> + 0.2% FeSO <sub>4</sub> at 15, 30 and 45 DAS	5.86	11.59
T <sub>5</sub>		50:25:25 kg NPK ha <sup>-1</sup>	5.74	11.26
T <sub>6</sub>		50:25:25 kg NPK ha <sup>-1</sup> + FS 0.5% ZnSO <sub>4</sub> + 0.2% FeSO <sub>4</sub> at 15, 30 and 45 DAS	5.94	11.85
T <sub>7</sub>	30 × 15 cm Spacing	40:20:0 kg NPK ha <sup>-1</sup> (RDF)	3.17	8.12
T <sub>8</sub>		40:20:0 kg NPK ha <sup>-1</sup> + FS 0.5% ZnSO <sub>4</sub> + 0.2% FeSO <sub>4</sub> at 15, 30 and 45 DAS	3.47	8.85
T <sub>9</sub>		40:20:20 kg NPK ha <sup>-1</sup>	4.15	9.37
T <sub>10</sub>		40:20:20 kg NPK ha <sup>-1</sup> + FS 0.5% ZnSO <sub>4</sub> + 0.2% FeSO <sub>4</sub> at 15, 30 and 45 DAS	4.89	10.59
T <sub>11</sub>		50:25:25 kg NPK ha <sup>-1</sup>	4.58	10.36
T <sub>12</sub>		50:25:25 kg NPK ha <sup>-1</sup> + FS 0.5% ZnSO <sub>4</sub> + 0.2% FeSO <sub>4</sub> at 15, 30 and 45 DAS	5.24	10.83
SEd			0.17	0.32
CD (p=0.05)			0.36	0.69



**Table 4. Effect of plant geometry, soil and foliar nutrition on crop growth rate (kg ha<sup>-1</sup>day<sup>-1</sup>) of rainfed dual purpose sorghum**

Treatments			Crop growth rate (kg ha <sup>-1</sup> day <sup>-1</sup> )		
			30 DAS	60 DAS	At harvest
T <sub>1</sub>	45 × 15 cm spacing	40:20:0 kg NPK ha <sup>-1</sup> (RDF)	38.26	211.48	154.26
T <sub>2</sub>		40:20:0 kg NPK ha <sup>-1</sup> + FS 0.5% ZnSO <sub>4</sub> + 0.2% FeSO <sub>4</sub> at 15, 30 and 45 DAS	43.17	213.84	159.94
T <sub>3</sub>		40:20:20 kg NPK ha <sup>-1</sup>	48.36	223.37	162.16
T <sub>4</sub>		40:20:20 kg NPK ha <sup>-1</sup> + FS 0.5% ZnSO <sub>4</sub> + 0.2% FeSO <sub>4</sub> at 15, 30 and 45 DAS	55.03	258.21	165.98
T <sub>5</sub>		50:25:25 kg NPK ha <sup>-1</sup>	52.13	241.42	162.36
T <sub>6</sub>		50:25:25 kg NPK ha <sup>-1</sup> + FS 0.5% ZnSO <sub>4</sub> + 0.2% FeSO <sub>4</sub> at 15, 30 and 45 DAS	56.25	266.79	167.99
T <sub>7</sub>	30 × 15 cm Spacing	40:20:0 kg NPK ha <sup>-1</sup> (RDF)	38.32	212.17	159.18
T <sub>8</sub>		40:20:0 kg NPK ha <sup>-1</sup> + FS 0.5% ZnSO <sub>4</sub> + 0.2% FeSO <sub>4</sub> at 15, 30 and 45 DAS	45.20	222.95	161.17
T <sub>9</sub>		40:20:20 kg NPK ha <sup>-1</sup>	58.11	270.15	169.55
T <sub>10</sub>		40:20:20 kg NPK ha <sup>-1</sup> + FS 0.5% ZnSO <sub>4</sub> + 0.2% FeSO <sub>4</sub> at 15, 30 and 45 DAS	62.38	297.41	181.81
T <sub>11</sub>		50:25:25 kg NPK ha <sup>-1</sup>	61.37	291.39	179.04
T <sub>12</sub>		50:25:25 kg NPK ha <sup>-1</sup> + FS 0.5% ZnSO <sub>4</sub> + 0.2% FeSO <sub>4</sub> at 15, 30 and 45 DAS	66.23	317.12	192.96
SEd			1.63	8.25	5.37
CD (p=0.05)			3.47	17.48	11.29



**Table 5. Effect of plant geometry, soil and foliar nutrition on grain yield and stover yield (kg ha<sup>-1</sup>) of rainfed dual purpose sorghum**

Treatments		Grain yield (kg ha <sup>-1</sup> )	Stover yield (kg ha <sup>-1</sup> )	
T <sub>1</sub>	45 × 15 cm spacing	40:20:0 kg NPK ha <sup>-1</sup> (RDF)	2572	10184
T <sub>2</sub>		40:20:0 kg NPK ha <sup>-1</sup> + FS 0.5% ZnSO <sub>4</sub> + 0.2% FeSO <sub>4</sub> at 15, 30 and 45 DAS	2745	10423
T <sub>3</sub>		40:20:20 kg NPK ha <sup>-1</sup>	2905	10792
T <sub>4</sub>		40:20:20 kg NPK ha <sup>-1</sup> + FS 0.5% ZnSO <sub>4</sub> + 0.2% FeSO <sub>4</sub> at 15, 30 and 45 DAS	3074	11896
T <sub>5</sub>		50:25:25 kg NPK ha <sup>-1</sup>	2942	11184
T <sub>6</sub>		50:25:25 kg NPK ha <sup>-1</sup> + FS 0.5% ZnSO <sub>4</sub> + 0.2% FeSO <sub>4</sub> at 15, 30 and 45 DAS	3205	11996
T <sub>7</sub>	30 × 15 cm spacing	40:20:0 kg NPK ha <sup>-1</sup> (RDF)	2743	10238
T <sub>8</sub>		40:20:0 kg NPK ha <sup>-1</sup> + FS 0.5% ZnSO <sub>4</sub> + 0.2% FeSO <sub>4</sub> at 15, 30 and 45 DAS	2803	10768
T <sub>9</sub>		40:20:20 kg NPK ha <sup>-1</sup>	3321	12032
T <sub>10</sub>		40:20:20 kg NPK ha <sup>-1</sup> + FS 0.5% ZnSO <sub>4</sub> + 0.2% FeSO <sub>4</sub> at 15, 30 and 45 DAS	3716	13130
T <sub>11</sub>		50:25:25 kg NPK ha <sup>-1</sup>	3562	12876
T <sub>12</sub>		50:25:25 kg NPK ha <sup>-1</sup> + FS 0.5% ZnSO <sub>4</sub> + 0.2% FeSO <sub>4</sub> at 15, 30 and 45 DAS	3961	13972
SEd		110.50	395.40	
CD (p=0.05)		236.00	840.76	